



ACADEMIC UROLOGY

Pottstown | Phoenixville

Robert M. Rose, M.D.
John E. Leech, Jr., M.D.
José G. Moreno, M.D.
Ronald L. Kabler, M.D.
Pankaj Kalra, M.D.
Lisa A. Robbins, CRNP
Julia C. Coyle, PA-C

MEDICATIONS TO STOP – IN OFFICE PROCEDURES

(Before stopping any medication, check with your personal physician or internist)

LOVENOX stop 24 hours prior to surgery.

ELIQUIS, PRADAXA & XARELTO stop 72 hours (3 days) prior to surgery.

COUMADIN stop 5 days prior to surgery.

PLAVIX (Clopidogrel Bisulfate) stop 7 days prior to surgery.

DO NOT TAKE THE FOLLOWING MEDICATIONS STARTING 10 DAYS PRIOR TO SURGERY:

Aspirin: (Anacin, Bayer, Bayer Children's Chewable Aspirin, Bayer Select, Backache, Bufferin, Ecotrin, Excedrin, Etc.)

Drugs that contain Aspirin or Aspirin-like compounds:

Aggrenox	COPE	Magnesium Salicylate
Alka Seltzer	Darvon	Midol
Anaprox	Doan's	Momentum
Arthritis Pain Formula	Dristan	Percodan
Ascripton	Duragesic	Sodium Salicylate
Aspergum	Empirin	Triaminicin
Aspirin Compound	Extra Strength Bayer	Vanquish
Aspirin Compound with Codeine	Fiorinal	

Non-Steroidal Anti-Inflammatory Agents:

Advil	Feldene (Piroxicam)	Orudis
Advil Cold and Sinus	Fenoprofen (Nalfon)	Relafen (Nabumetone)
Aleve	Haltran	Rufen
Arthritis Foundation IB Tabs	Ibuprofen	Tolectin (Tolmetin Sodium)
Asacol	Indomethacin (Indocin)	Toradol (Ketorolac Trimethamine)
Bayer Select IB Pain Relief	Lodine (Etodolac)	Trilistate
Cataflam (diclofenac potassium)	Medipren	Voltarin (Diclofenac Sodium)
Celebrex	Mobic	
Clinoril (Salindac)	Motrin IB	
Cramp End Tablets	Nalfon (Fenoprofen)	
Dolobid (Diflunisal)	Naprosyn (Naproxen Sodium)	
Excedrin IB	Nuprin	

Other Medications that may cause bleeding:

Dicumarol	Persantine (Dipyridamole)	Trental (Pentoxifylline)
Heparin	Phenylbutazone	Multi-Vitamin/Vitamin E/Fish Oil
Oxyphenbutazone (Oxalid, Tandearil)	Ticlid (ticlopidine hydrochloride)	Over the Counter Supplements

You **May** take Tylenol, any acetaminophen containing product or narcotics such as Codeine, Tylenol #3, Tylox, Extra-strength Tylenol with Codeine, Aspirin Free Anacin, Fioricet, Percocet, Vicodin, Contac, Sine-Aid.

Fenfluramin (Pondimin) should be discontinued gradually one week prior to any elective surgery. Please check with your family physician regarding instructions on how to gradually stop this medication.

11/18/2015



ACADEMIC UROLOGY

Pottstown | Phoenixville

Robert M. Rose, M.D.
John E. Leech, Jr., M.D.
José G. Moreno, M.D.
Ronald L. Kabler, M.D.
Pankaj Kalra, M.D.
Lisa A. Robbins, CRNP
Julia C. Coyle, PA-C

MEDICATIONS TO STOP

(Before stopping any medication, check with your personal physician or internist)

GLUCOPHAGE (Metformin) and GLUCOVANCE stop 24 hrs prior to surgery.
LOVENOX stop 24 hours prior to surgery.
ELIQUIS, PRADAXA & XARELTO stop 72 hours (3 days) prior to surgery.
COUMADIN stop 5 days prior to surgery.
PLAVIX (Clopidogrel Bisulfate) stop 7 days prior to surgery.

DO NOT TAKE THE FOLLOWING MEDICATIONS STARTING 10 DAYS PRIOR TO SURGERY:

Aspirin: (Anacin, Bayer, Bayer Children's Chewable Aspirin, Bayer Select, Backache, Bufferin, Ecotrin, Excedrin, Etc.)

Drugs that contain Aspirin or Aspirin-like compounds:

Aggrenox	COPE	Magnesium Salicylate
Alka Seltzer	Darvon	Midol
Anaprox	Doan's	Momentum
Arthritis Pain Formula	Dristan	Percodan
Ascripton	Duragesic	Sodium Salicylate
Aspergum	Empirin	Triaminicin
Aspirin Compound	Extra Strength Bayer	Vanquish
Aspirin Compound with Codeine	Fiorinal	

Non-Steroidal Anti-Inflammatory Agents:

Advil	Feldene (Piroxicam)	Orudis
Advil Cold and Sinus	Fenoprofen (Nalfon)	Relafen (Nabumetone)
Aleve	Haltran	Rufen
Arthritis Foundation IB Tabs	Ibuprofen	Tolectin (Tolmetin Sodium)
Asacol	Indomethacin (Indocin)	Toradol (Ketorolac Trimethamine)
Bayer Select IB Pain Relief	Lodine (Etodolac)	Trilistate
Cataflam (diclofenac potassium)	Medipren	Voltarin (Diclofenac Sodium)
Celebrex	Mobic	
Clinoril (Salindac)	Motrin IB	
Cramp End Tablets	Nalfon (Fenoprofen)	
Dolobid (Diflunisal)	Naprosyn (Naproxen Sodium)	
Excedrin IB	Nuprin	

Other Medications that may cause bleeding:

Dicumarol	Persantine (Dipyridamole)	Trental (Pentoxifylline)
Heparin	Phenylbutazone	Multi-Vitamin/Vitamin E/Fish Oil
Oxyphenbutazone (Oxalid, Tandearil)	Ticlid (ticlopidine hydrochloride)	Over the Counter Supplements

You **May** take Tylenol, any acetaminophen containing product or narcotics such as Codeine, Tylenol #3, Tylox, Extra-strength Tylenol with Codeine, Aspirin Free Anacin, Fioricet, Percocet, Vicodin, Contac, Sine-Aid.

Fenfluramin (Pondimin) should be discontinued gradually one week prior to any elective surgery. Please check with your family physician regarding instructions on how to gradually stop this medication.

11/18/2015

BETA-BLOCKER MEDICATIONS

New Government Rule states that a patient must take all beta-blocker medications prescribed with a sip of water in the morning prior to surgery.

Atenolol
Atenolo/chlorthalidone
Betapace
Betapace AF
Betaxolol
Bisoprolol
Bisoprolol/fumarate
Bisoprolol/hydrochlorothiazide
Blocadren
Brevibloc
Bystolic
Carteolol
Cartrol
Carvedilol
Coreg
Corgard
Corzide 40/5
Corzide 80/5
Esmolol
Inderal
Inderal LA
Inderide
Inderide LA
InnoPran XL
Kerlone
Labetalol
Levatol
Lopressor
Lopressor HCT
Lopressor/hydrochlorothiazide
Metoprolol
Metoprolol/hydrochlorothiazide
Metoprolol Tartrate/hydrochlorothiazide
Nadolol
Nadolol/bendroflumethiazide
Nebivolol
Nebivolol HCL
Nebivolol Hydrochloride
Nomadyne
Penbutolol
Pindolol
Propranolol
Propranolol HCL

Propranolol Hydrochloride
Propranolol/hydrochlorothiazide
Sectral
Sorine
Sotolol
Sotalol HCL
Tenoretic
Tenormin
Tenormin I.V.
Timolide
Timolol
Timolol Maleate/hydrochlorothiazide
Timolol/hydrochlorothiazide
Toprol
Toprol-XL
Trandate
Trandate HCL
Visken
Zebeta
Ziac